

Anisotropic Non-gaussianity with noncommutative spacetime

Akhilesh Nautiyal

The Institute of Mathematical Sciences, Taramani, Chennai, India

(Dated: March 19, 2013)

Abstract

We study single field inflation in noncommutative spacetime and compute two-point and three-point correlation functions for the curvature perturbation. We find that both power spectrum and bispectrum for comoving curvature perturbation are statistically anisotropic and the bispectrum is also modified by a phase factor depending upon the noncommutative parameters. The non-linearity parameter f_{NL} is small for small statistical anisotropic corrections to the bispectrum coming from the noncommutative geometry. There is a scale dependence of f_{NL} due to the noncommutative spacetime which is different from the standard single field inflation models and statistically anisotropic vector field inflation models. The anisotropy of the two-point function and the running of the f_{NL} parameter will be tested precisely from the data which will be released soon from the PLANCK experiment.

PACS numbers: 98.80.Cq, 11.10.Nx, 98.70.Vc

1. INTRODUCTION

Inflation [1] not only solves the various puzzles of the Big-Bang theory, but it also provides seeds for the temperature anisotropy of the cosmic microwave background (CMB) radiation and structures in the universe. In the standard inflationary scenario, the potential energy of a scalar field called "inflaton" dominates the energy density of the universe and its quantum fluctuations generate perturbations in the metric causing small inhomogeneities in the early universe which give rise to CMB anisotropy and structures in the universe.

Inflation predicts nearly scale invariant, adiabatic and gaussian perturbations. The first two are in excellent agreement with the observations of CMB anisotropy and polarization by COBE [2], WMAP [3–5] and other ground based and satellite based experiments, but the test of gaussian statistics of the perturbations is controversial and is the major goal of ongoing and future observations like Planck [6], CMBPOL [7] and Euclid satellite [8].

The non-gaussianity in CMB can be primordial or can be generated due to secondary sources (see [9] for detailed review). The primordial non-gaussianity arises due to the interaction terms in the scalar potential and non-linearities of the gravity, where the latter effect is dominant than the former. The magnitude of the non-gaussianity in standard single-field inflation comes out to be small and of the order of slow-roll parameters [10].

Inflation occurs at a very high energy and it stretches out very small scales, of the order of Planck length, to the current hubble scale due to superluminal expansion of the universe. So it provides window to see the new physics at the Planck scale at which quantum corrections to the gravity becomes important. These new effects can give rise to large primordial non-gaussianity that can be seen by Planck.

Spacetime noncommutativity (see [11] for review) is one of such modifications at high-energy, which is well motivated by quantum gravity and string theory. Modifications to the power spectrum of scalar perturbations during inflation and its effects on CMB due to noncommutative geometry has been studied in many places [12–15]. In this paper we compute the three-point correlation functions of the curvature perturbation and hence the non-linearity parameter f_{NL} determining the primordial non-gaussianity using the noncommutative quantum field theories related to deformed Poincare symmetry [12]. In this approach of noncommutative geometry quantum fields follow twisted statistics, as implied by the deformed Poincare symmetry in quantum theories. Non-gaussianity due to noncommutative spacetime of [12] has also been calculated in [16] using the δN formalism [17], which is used to calculate the local non-gaussianity and treats the comoving curvature perturbations as classical. But, here we compute the two-point and three-point function using Maldacena approach [10] which is based on the second order perturbation theory and is valid for all kinds of non-gaussian effects.

2. QUANTUM FIELDS IN NONCOMMUTATIVE SPACETIME

At the energy scale of inflation, the noncommutativity of spacetime, which is motivated by Heisenberg uncertainty principle and Einstein's general relativity, can play a crucial role. The spacetime noncommutativity can be represented by the commutation relations [11]

$$[\tilde{x}_\mu, \tilde{x}_\nu] = i\theta_{\mu\nu} \tag{1}$$

where $\theta_{\mu\nu}$ is a real antisymmetric matrix with constant elements and $\tilde{x}_\mu$ are the coordinate functions of the chosen coordinate system:

$$\tilde{x}_\mu(x) = x_\mu. \quad (2)$$

The relation (1) holds only in special coordinate systems and looks quite complicated in other coordinates. The natural choice of the coordinate for cosmological applications is the comoving frame, where the galaxies are freely falling. We chose $\theta_{\mu\nu}$ to be constant in the comoving coordinates as it makes our computation convenient and the physical scales can be computed by using the relations

$$\theta_{0i}^{ph}(t) = a(t)\theta_{0i}, \quad \theta_{ij}^{ph}(t) = a^2(t)\theta_{ij} \quad (3)$$

Our choice is opposite to the choice of [13] where they assume $\theta_{\mu\nu}^{ph}$ to be constant in time.

Due to spacetime noncommutativity, the usual quantum fields are deformed and can be given in terms of undeformed quantum fields as [11]

$$\phi_\theta = \phi_0 e^{\frac{1}{2} \overleftarrow{\partial} \wedge P} \quad (4)$$

where $\overleftarrow{\partial} \wedge P = \overleftarrow{\partial}_\mu \theta^{\mu\nu} P_\nu$ and P_ν represents the field momentum operator. The product of the deformed (twisted) quantum fields at the same spacetime point is represented by the star product given as

$$(\phi_\theta \star \phi_\theta)(x) = \phi_\theta(x) e^{\frac{i}{2} \overleftarrow{\partial}_x \wedge \overrightarrow{\partial}_y} \phi_\theta(y) \Big|_{x=y} \quad (5)$$

In the following sections we will make use of these relations to calculate two and three-point correlation functions of the comoving curvature perturbations.

3. TWO-POINT AND THREE-POINT FUNCTION WITH NONCOMMUTATIVE SPACETIME

3.1. Background

The action of a single scalar field minimally coupled with gravity is

$$S = \int d^4x \sqrt{-g} \left(\frac{M_p^2}{2} R + \mathcal{L} \right). \quad (6)$$

Here R is the Ricci scalar and $\mathcal{L}$ is the Lagrangian for the scalar field i.e.

$$\mathcal{L} = -\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi). \quad (7)$$

Noncommutativity doesn't change the classical background so all the background dynamics will be similar to the standard case. We take the metric signature $(-, +, +, +)$ and work in the units where $M_p = 1$. The background geometry of the homogeneous isotropic universe is described by the FRW metric

$$ds^2 = -dt^2 + a^2(t) (dx^2 + dy^2 + dz^2). \quad (8)$$

For a scalar field dominated universe the Friedmann equations are given as

$$\begin{aligned} 3H^2 &= \frac{1}{2}\dot{\phi}^2 + V(\phi) \\ \dot{\rho} + 3H(\rho + p) &= 0 \\ \dot{H} &= -\frac{1}{2}(\rho + p) \end{aligned} \quad (9)$$

The equation of motion for the scalar field is given as

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV(\phi)}{d\phi} = 0. \quad (10)$$

During inflation, the potential energy of the scalar field dominates the total energy density of the universe and the dynamics of the scalar field is governed by slow-roll parameters defined as as

$$\begin{aligned} \epsilon &= -\frac{\dot{H}}{H^2} \\ \eta &= \frac{\dot{\epsilon}}{\epsilon H} \end{aligned} \quad (11)$$

Here we follow the definition of η as in [18], which is different from the definition using scalar field potential ($\eta_V = \frac{d^2V}{d\phi^2}$) and $\eta = -2\eta_V + 4\epsilon$.

3.2. Perturbations and ADM formalism

The quantum fluctuations in the scalar field $\delta\phi(x, t)$ generated during inflation are coupled to the perturbations in the metric through Einstein's equation. Inflation gives rise to scalar and tensor perturbations in the metric and the scalar part is written as

$$ds^2 = -(1 + 2\Phi)dt^2 + 2a^2(t)B_{,i}dx^i dt + a^2(t)((1 - 2\Psi)\delta_{ij} + 2E_{,ij})dx^i dx^j \quad (12)$$

Here we have four scalar degrees of freedom in the metric and one in the scalar field which can be reduced to three by using gauge transformations. We can again use the constraint equations derived from the perturbed Einstein's equation and describe the scalar perturbations in terms of the curvature perturbation defined as [19]

$$\zeta = -\Psi - \frac{H}{\dot{\phi}}\delta\phi. \quad (13)$$

This variable is gauge invariant and is conserved on super-horizon scales. One can write the action (6) in terms of ζ and it turns out quadratic in ζ . To do the perturbation theory in higher order it is convenient to use ADM formalism where the metric can be written as [20]

$$ds^2 = -N^2 dt^2 + h_{ij}(dx^i + N^i dt)(dx^j + N^j dt) \quad (14)$$

where N is laps function, N_i, N_j are shift vectors and h_{ij} is metric of three-dimensional hypersurface of constant time. Here N and N_i appear as Lagrangian multipliers in the action so one can solve there constraint equation of motion and substitute the solution back into the action. This simplifies the tedious calculations needed while working with (12). Now we chose comoving gauge $\delta\phi = 0$ to do our calculation and in this gauge we can use non-linear generalization of ζ [21] and define the gauge as [10, 18]

$$h_{ij} = a^2 e^{2\zeta} \delta_{ij}, \quad \delta\phi = 0. \quad (15)$$

With this gauge choice the action (6) with the metric (14) becomes

$$S = \frac{1}{2} \int dt d^3x \sqrt{h} \left(N R^{(3)} - 2NV(\phi) + N^{-1} \dot{\phi}^2 + N^{-1} (E_{ij} E^{ij} - E^2) \right) \quad (16)$$

Here $R^{(3)}$ represents the Ricci scalar calculated using the three-dimensional metric h_{ij} and E_{ij} is related to the extrinsic curvature of the constant time hypersurface and is given as

$$E_{ij} = \frac{1}{2} \left(\dot{h}_{ij} - \nabla_j N_i - \nabla_i N_j \right). \quad (17)$$

Varying the action (16) we get the constraint equation for N and N^i as

$$\begin{aligned} R^{(3)} - 2V - N^{-2} (E_{ij} E^{ij} - E^2) - N^{-2} \dot{\phi}^2 &= 0, \\ \nabla_j [N^{-1} (E_i^j - \delta_i^j E)] &= 0. \end{aligned} \quad (18)$$

Now we can decompose N_i into irrotational and incompressible parts as $N_i = \tilde{N}_i + \partial_i \psi$ where $\partial_i \tilde{N}^i = 0$ and expand N, ψ and $\tilde{N}^i$ into powers of ζ as

$$\begin{aligned} N &= 1 + \alpha_1 + \alpha_2 + \dots \\ \tilde{N}_i &= \tilde{N}_i^{(1)} + \tilde{N}_i^{(2)} + \dots \\ \psi &= \psi_1 + \psi_2 + \dots \end{aligned} \quad (19)$$

Using these expansions, the constraint equations (18) can be solved order by order with metric (15) and at first order one gets

$$\alpha_1 = \frac{\dot{\zeta}}{H}, \quad \tilde{N}_i^{(1)} = 0, \quad \psi_1 = -\frac{\zeta}{H} + \chi, \quad \partial^2 \chi = a^2 \epsilon \dot{\zeta} \quad (20)$$

Here $\partial^2 = \delta^{ij} \partial_i \partial_j$ and the use of suitable choice of boundary conditions has been made to put $N_i^{(1)} = 0$. As mentioned in [10, 18] to calculate the action up to n^{th} order in ζ , we need to calculate N and N_i only up to the order- ζ^{n-1} and here the terms of order- ζ^2 also drop out from the third order action, so equation (20) is sufficient to compute the action up to third order. So, after putting these solutions in (16) we get the action for second and third order in ζ as [10, 18, 22]

$$S_2 = \int dt d^3x \left[a^3 \epsilon \dot{\zeta}^2 - a \epsilon (\partial \zeta)^2 \right] \quad (21)$$

$$\begin{aligned} S_3 = \int dt d^3x & \left[-a \epsilon \zeta (\partial \zeta)^2 - a^3 \epsilon \dot{\zeta}^3 + 3a^3 \epsilon \zeta \dot{\zeta}^2 \right. \\ & \left. + \frac{1}{2a} \left(3\zeta - \frac{\dot{\zeta}}{H} \right) (\partial_i \partial_j \psi \partial^i \partial^j \psi - \partial^2 \psi \partial^2 \psi) - 2a^{-1} \partial_i \psi \partial_i \zeta \partial^2 \psi \right] \end{aligned} \quad (22)$$

3.3. Two-point correlation function and power spectrum

Now to calculate the two-point correlation function the quadratic part (21) of the action is considered, which in conformal time ($d\tau = \frac{dt}{a}$) can be written as

$$S_2 = \int d\tau d^3x a^2 \epsilon [\zeta'^2 - (\partial\zeta)^2] \quad (23)$$

Here $\prime$ denotes derivative w.r.t conformal time τ . The above action looks like an action of a massless scalar field in conformal spacetime and ζ can be considered as the scalar field for quantization. ζ can be written in terms of creation and annihilation operator as

$$\zeta(\vec{x}, \tau) = \int \frac{d^3k}{(2\pi)^3} \zeta(\vec{k}, \tau) e^{i\vec{k}\cdot\vec{x}} = \int \frac{d^3k}{(2\pi)^3} \left(u(\vec{k}, \tau) a_{\vec{k}} + u^*(-\vec{k}, \tau) a_{-\vec{k}}^\dagger \right) e^{i\vec{k}\cdot\vec{x}} \quad (24)$$

The equation of motion for ζ can be obtained by varying the action (23) and is given by

$$\zeta'' + 2\frac{z'}{z}\zeta' - \partial^2\zeta = 0. \quad (25)$$

Here $z^2 = 2a^2\epsilon$ and we can define $v_{\vec{k}} = z\zeta(\vec{k}, \tau)$ and use equation (24) to get

$$v_{\vec{k}}'' + \left(k^2 - \frac{z''}{z} \right) v_{\vec{k}} = 0. \quad (26)$$

The solution for the mode functions $v_{\vec{k}}$ can be obtained assuming Bunch Davies initial conditions and is given as

$$v_{\vec{k}} = \frac{1}{\sqrt{2k}} \left(1 - \frac{i}{k\tau} \right) e^{-ik\tau}. \quad (27)$$

Hence the basis function $u(\vec{k}, \tau)$ is

$$u(\vec{k}, \tau) = \frac{v_{\vec{k}}}{z} = \frac{iH}{\sqrt{4\epsilon k^3}} (1 + ik\tau) e^{-ik\tau}. \quad (28)$$

The two point correlation function of the field ζ in position space can be expressed as

$$\langle \zeta(\vec{x}, \tau) \zeta(\vec{y}, \tau) \rangle = \int \frac{d^3k d^3k'}{(2\pi)^6} \langle 0 | \zeta(\vec{k}, \tau) \zeta(\vec{k}', \tau) | 0 \rangle e^{i(\vec{k}\cdot\vec{x} + \vec{k}'\cdot\vec{y})} \quad (29)$$

$$= \int \frac{d^3k}{(2\pi)^3} |u(\vec{k}, \tau)|^2 e^{i\vec{k}\cdot(\vec{x}-\vec{y})}, \quad (30)$$

where we have used the relation $\langle 0 | \zeta(\vec{k}, \tau) \zeta(\vec{k}', \tau) | 0 \rangle = (2\pi)^3 \delta^3(\vec{k} + \vec{k}') u(\vec{k}, \tau) u^*(-\vec{k}', \tau)$.

The power spectrum for ζ is defined by

$$\langle 0 | \zeta(\vec{k}, \tau) \zeta(\vec{k}', \tau) | 0 \rangle = (2\pi)^3 \delta^3(\vec{k} + \vec{k}') P_\zeta(k) \quad (31)$$

So

$$P_\zeta(k) = |u(\vec{k}, \tau)|^2 \quad (32)$$

The another convention for the power spectrum, that is commonly used for data analysis, is

$$\Delta_\zeta^2 = \frac{k^3}{2\pi^2} |u(\vec{k}, \tau)|^2 \quad (33)$$

In this case Δ_ζ represents the variance of the classical fluctuations and the two-point correlation in position space becomes

$$\langle \zeta(\vec{x}, \tau) \zeta(\vec{y}, \tau) \rangle = \int \frac{dk}{k} \Delta_\zeta^2 e^{i\vec{k} \cdot (\vec{x} - \vec{y})}. \quad (34)$$

The power spectrum is calculated on super-horizon limit i.e. $-k\tau \ll 1$ in which $v_{\vec{k}} = \frac{1}{\sqrt{2k}} \left(-\frac{i}{k\tau}\right) e^{-ik\tau}$ and we get the power spectrum as

$$P_\zeta(k) = \frac{H^2}{4\epsilon} \frac{1}{k^3} \quad (35)$$

Now due to noncommutativity of spacetime the two point correlation function for field ζ gets modified [12]. We will denote the field in noncommutative spacetime with a subscript θ . Since here ζ represents our quantum field, hence similar to (4) the twisted quantum field ζ_θ can be expressed in terms of the untwisted field ζ as

$$\zeta_\theta(\vec{x}, t) = \zeta(\vec{x}, t) e^{\frac{1}{2} \overleftarrow{\partial}_\mu \wedge P_\nu}. \quad (36)$$

With the twisted quantum field one can compute the two-point correlation function in position space as

$$\begin{aligned} \langle \zeta_\theta(\vec{x}, t) \zeta_\theta(\vec{y}, t') \rangle &= \langle \zeta(\vec{x}, t) e^{\frac{1}{2} \overleftarrow{\partial}_{x_\mu} \wedge P_\nu} \zeta(\vec{y}, t') e^{\frac{1}{2} \overleftarrow{\partial}_{y_\mu} \wedge P_\nu} \rangle \\ &= \langle \zeta(\vec{x}, t) \zeta(\vec{y}, t') \rangle e^{-\frac{i}{2} \overleftarrow{\partial}_{x_\mu} \wedge \overrightarrow{\partial}_{y_\nu}}, \end{aligned} \quad (37)$$

where we have used the commutation relations between the field and the momentum operator $[P_\mu, \zeta] = -i\partial_\mu \zeta$. Now taking the Fourier transform on the right hand side we get

$$\begin{aligned} \langle \zeta_\theta(\vec{x}, t) \zeta_\theta(\vec{y}, t') \rangle &= \int \frac{d^3k d^3k'}{(2\pi)^6} \langle 0 | \zeta(\vec{k}, t) \zeta(\vec{k}', t') | 0 \rangle e^{-\frac{i}{2} \overleftarrow{\partial}_{x_\mu} \wedge \overrightarrow{\partial}_{y_\nu}} e^{i(\vec{k} \cdot \vec{x} + \vec{k}' \cdot \vec{y})} \\ &= \int \frac{d^3k d^3k'}{(2\pi)^6} \langle 0 | \zeta(\vec{k}, t) \zeta(\vec{k}', t') | 0 \rangle e^{-\frac{i}{2} (\partial_t \theta^{0i} \partial_{\vec{y}} + \partial_{\vec{x}} \theta^{i0} \partial_{t'} + \partial_{\vec{x}} \wedge \partial_{\vec{y}})} e^{i(\vec{k} \cdot \vec{x} + \vec{k}' \cdot \vec{y})} \\ &= \int \frac{d^3k d^3k'}{(2\pi)^6} \langle 0 | \zeta(\vec{k}, t) \zeta(\vec{k}', t') | 0 \rangle e^{\left(\frac{i}{2} \vec{k} \wedge \vec{k}' + \frac{\vec{\theta}^0 \cdot \vec{k}'}{2} \partial_t - \frac{\vec{\theta}^0 \cdot \vec{k}}{2} \partial_{t'}\right)} e^{i(\vec{k} \cdot \vec{x} + \vec{k}' \cdot \vec{y})} \\ &= \int \frac{d^3k d^3k'}{(2\pi)^6} \langle 0 | \zeta \left(\vec{k}, t + \frac{\vec{\theta}^0 \cdot \vec{k}'}{2} \right) \zeta \left(\vec{k}', t' - \frac{\vec{\theta}^0 \cdot \vec{k}}{2} \right) | 0 \rangle e^{\frac{i}{2} \vec{k} \wedge \vec{k}'} e^{i(\vec{k} \cdot \vec{x} + \vec{k}' \cdot \vec{y})} \end{aligned} \quad (38)$$

Here $\vec{\theta}^0 = \theta^{0i}$. So the two-point correlation function in momentum space can be expressed as

$$\langle 0 | \zeta(\vec{k}, t) \zeta(\vec{k}', t') | 0 \rangle = e^{\frac{i}{2} \vec{k} \wedge \vec{k}'} \langle 0 | \zeta \left(\vec{k}, t + \frac{\vec{\theta}^0 \cdot \vec{k}'}{2} \right) \zeta \left(\vec{k}', t' - \frac{\vec{\theta}^0 \cdot \vec{k}}{2} \right) | 0 \rangle \quad (39)$$

Now since in de Sitter space

$$\tau(t) = \frac{1}{aH} e^{-Ht} \quad (40)$$

So in conformal time and in the limit $t' \rightarrow t$

$$\zeta \left(\vec{k}, t + \frac{\vec{\theta}^0 \cdot \vec{k}'}{2} \right) \rightarrow \zeta \left(\vec{k}, \tau e^{-H \frac{\vec{\theta}^0 \cdot \vec{k}'}{2}} \right) \quad (41)$$

$$\zeta \left(\vec{k}, t' - \frac{\vec{\theta}^0 \cdot \vec{k}}{2} \right) \rightarrow \zeta \left(\vec{k}, \tau e^{H \frac{\vec{\theta}^0 \cdot \vec{k}}{2}} \right) \quad (42)$$

Hence the two-point function will be

$$\begin{aligned} \langle \zeta(\vec{k}, \tau) \zeta(\vec{k}', \tau) \rangle &= \langle 0 | \zeta \left(\vec{k}, \tau e^{-H \frac{\vec{\theta}^0 \cdot \vec{k}'}{2}} \right) \zeta \left(\vec{k}', \tau e^{H \frac{\vec{\theta}^0 \cdot \vec{k}}{2}} \right) | 0 \rangle e^{\frac{i}{2} \vec{k} \wedge \vec{k}'} \\ &= \left| u \left(\vec{k}, \tau e^{H \frac{\vec{\theta}^0 \cdot \vec{k}}{2}} \right) \right|^2 (2\pi)^3 \delta^3(\vec{k} + \vec{k}') \end{aligned} \quad (43)$$

Now we take the self-adjoint part of two-point correlation function defined as [12]

$$\langle 0 | \zeta_\theta(\vec{k}, \tau) \zeta_\theta(\vec{k}', \tau) | 0 \rangle_M = \frac{1}{2} \left(\langle 0 | \zeta_\theta(\vec{k}, \tau) \zeta_\theta(\vec{k}', \tau) | 0 \rangle + \langle 0 | \zeta_\theta(-\vec{k}, \tau) \zeta_\theta(-\vec{k}', \tau) \rangle \right) \quad (44)$$

So the power spectrum can be obtained from (31) as

$$P_{\zeta_\theta}(k) = \frac{1}{2} \left(\left| u \left(\vec{k}, \tau e^{H \frac{\vec{\theta}^0 \cdot \vec{k}}{2}} \right) \right|^2 + \left| u \left(-\vec{k}, \tau e^{-H \frac{\vec{\theta}^0 \cdot \vec{k}}{2}} \right) \right|^2 \right) \quad (45)$$

Since on super-horizon limit $v_{\vec{k}} = \frac{1}{\sqrt{2k}} \left(\frac{-i}{k\tau} e^{-H \frac{\vec{\theta}^0 \cdot \vec{k}}{2}} \right)$ so

$$P_{\zeta_\theta}(k) = P_\zeta(k) \cosh \left(H \vec{\theta}^0 \cdot \vec{k} \right) \quad (46)$$

This power spectrum was derived in [12, 16] and was tested with the CMB observations in [13] and it was found that the extra θ -dependent factor in the power spectrum can lead to direction dependent CMB anisotropy and future data at large l can tightly constraint the scale of noncommutativity.

3.4. Three-point function

The primordial non-gaussianity in CMB arises due to the non-zero three-point and four-point correlation functions of curvature perturbations. These correlation functions were calculated for noncommutative spacetime in [16], where they have used δN formalism which ignores modifications to the correlation functions at Hubble crossing and also interaction between quantum fluctuations on sub-hubble scales with the super-hubble scale fluctuations at non-linear label.

The third order action (22) obtained for ζ using ADM formalism is

$$S_3 = \int dt d^3x \left[-a\epsilon\zeta(\partial\zeta)^2 - a^3\epsilon\dot{\zeta}^3 + 3a^3\epsilon\zeta\dot{\zeta}^2 + \frac{1}{2a} \left(3\zeta - \frac{\dot{\zeta}}{H} \right) (\partial_i\partial_j\psi\partial^i\partial^j\psi - \partial^2\psi\partial^2\psi) - 2a^{-1}\partial_i\psi\partial_i\zeta\partial^2\psi \right] \quad (47)$$

We put the value of ψ from Eq. (20) in this action, integrate by parts and use background Friedmann equations to get terms proportional to ϵ^2

$$S_3 = \int dt d^3x \left[a^3\epsilon^2\zeta\dot{\zeta}^2 + a\epsilon^2\zeta(\partial\zeta)^2 - 2a\epsilon\dot{\zeta}(\partial\zeta)(\partial\chi) + \frac{a^3\epsilon}{2}\frac{d\eta}{dt}\zeta^2\dot{\zeta} + \frac{\epsilon}{2a}(\partial\zeta)(\partial\chi)(\partial^2\chi) + \frac{\epsilon}{4a}(\partial^2\zeta)(\partial\chi)^2 + \frac{1}{2}a\mathcal{F}\frac{\delta L}{\delta\zeta}\Big|_1 \right] \quad (48)$$

where $\mathcal{F} = (\eta\zeta^2 + \text{terms with derivatives of } \zeta)$ and $\frac{\delta L}{\delta\zeta}$ represents the terms proportional to the Gaussian action S_2 . We can again integrate by parts the above action to remove the terms involving $\partial\chi$ and use the Gaussian field equation (25) to get

$$S_3 = \int dt d^3x \left[4a^5\epsilon^2 H\dot{\zeta}^2\partial^{-2}\dot{\zeta} + \frac{1}{2}a\mathcal{F}\frac{\delta L}{\delta\zeta}\Big|_1 \right] \quad (49)$$

Now $\mathcal{F} = (\eta - \epsilon)\zeta^2 + 2\epsilon\partial^{-2}(\zeta\partial^2\zeta)$ and ∂^{-2} is the inverse of ∂^2 and we have ignored the terms containing the derivatives of ζ in $\mathcal{F}$ as they are negligible on super-horizon scales. One can get rid of the second term in the above action following field redefinition

$$\zeta \rightarrow \zeta_n + \frac{\mathcal{F}}{4}\zeta_n^2 \quad (50)$$

After this field redefinition the three-point function becomes

$$\begin{aligned} \langle \zeta(x_1)\zeta(x_2)\zeta(x_3) \rangle &= \langle \zeta_n(x_1)\zeta_n(x_2)\zeta_n(x_3) \rangle + \frac{(\eta - \epsilon)}{4} (\langle \zeta_n(x_1)\zeta_n(x_2) \rangle \langle \zeta_n(x_1)\zeta_n(x_3) \rangle + \text{permutations}) \\ &+ \frac{\epsilon}{2}\partial_{x_1}^{-2} (\langle \zeta(x_1)\zeta(x_2) \rangle \partial_{x_1}^2 \langle \zeta_n(x_1)\zeta_n(x_3) \rangle + \text{permutations}) . \end{aligned} \quad (51)$$

The first term in above expression represents the three point function while the last two terms represents the corrections to the three point function due to field redefinition. We

will omit the subscript n in the following calculations. Now the interaction Hamiltonian to calculate the three-point function can be obtained from action (49) i.e

$$\mathcal{H}(t') = - \int d^3x 4a^5 \epsilon^2 H \dot{\zeta}^2 \partial^{-2} \dot{\zeta} \quad (52)$$

As mentioned earlier, we use ζ as the quantum field to compute the various correlation functions. Hence, to see the effects of noncommutative geometry on three point correlation function we replace the usual quantum field ζ with the twisted field ζ_θ both in the interaction Hamiltonian and in (51). Since the product of the twisted fields at the same spacetime point is given by the star-product [11], the interaction Hamiltonian will be given as

$$\mathcal{H}(t') = - \int d^3x 4a^5 \epsilon^2 H \dot{\zeta}_\theta \star \dot{\zeta}_\theta \star \partial^{-2} \dot{\zeta}_\theta \quad (53)$$

Using the relation (4) between the twisted and untwisted quantum field and expression for star product (5), the interaction Hamiltonian becomes

$$\mathcal{H}(t') = - \int d^3x 4a^5 \epsilon^2 H \dot{\zeta}^2 \partial^{-2} \dot{\zeta} e^{\frac{1}{2} \overleftarrow{\partial}_\mu \wedge P_\nu} \quad (54)$$

The first term in the RHS of (51) is computed using the in-in formalism [23] and is given by

$$\begin{aligned} \langle \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) \rangle &= -i \int_{t_0}^t dt' \langle 0 | [\zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3), \mathcal{H}(t')] | 0 \rangle \\ &= -i \int_{t_0}^t dt' (\langle 0 | \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) \mathcal{H}(t') | 0 \rangle - \langle 0 | \mathcal{H}(t') \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) | 0 \rangle) \end{aligned} \quad (55)$$

Here the three point function is calculated at equal time i.e $t_1 = t_2 = t_3 = t$. Initially we will write them differently for simplification but will put them equal before integration w.r.t t' . Let us now consider the first term of above equation with (54) and name it as (a). So

$$(a) = 4i\epsilon^2 \int dt' a^5 H \int d^3x \langle 0 | \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) \dot{\zeta}^2 \partial^{-2} \dot{\zeta} \Big|_{t', \vec{x}} e^{\frac{1}{2} \overleftarrow{\partial}_\mu \wedge P_\nu} | 0 \rangle \quad (56)$$

Using the relation (36) we can replace the twisted quantum fields in terms of the untwisted quantum fields and it gives

$$\begin{aligned} (a) &= 4i\epsilon^2 \int dt' a^5 H \int d^3x \langle 0 | \zeta(x_1) \zeta(x_2) \zeta(x_3) e^{-\frac{i}{2} (\overleftarrow{\partial}_{x_1} \wedge \overleftarrow{\partial}_{x_2} + \overleftarrow{\partial}_{x_2} \wedge \overleftarrow{\partial}_{x_3} + \overleftarrow{\partial}_{x_1} \wedge \overleftarrow{\partial}_{x_3})} \\ &\quad \times e^{\frac{1}{2} \overleftarrow{\partial}_{x_1} \wedge P} e^{\frac{1}{2} \overleftarrow{\partial}_{x_2} \wedge P} e^{\frac{1}{2} \overleftarrow{\partial}_{x_3} \wedge P} \dot{\zeta}^2 \partial^{-2} \dot{\zeta} \Big|_{t', \vec{x}} e^{\frac{1}{2} \overleftarrow{\partial}_x \wedge P} | 0 \rangle \\ &= 4i\epsilon^2 \int dt' a^5 H \int d^3x \langle 0 | \zeta(x_1) \zeta(x_2) \zeta(x_3) e^{-\frac{i}{2} (\overleftarrow{\partial}_{x_1} \wedge \overleftarrow{\partial}_{x_2} + \overleftarrow{\partial}_{x_2} \wedge \overleftarrow{\partial}_{x_3} + \overleftarrow{\partial}_{x_1} \wedge \overleftarrow{\partial}_{x_3})} \\ &\quad \times e^{-\frac{i}{2} (\overleftarrow{\partial}_{x_1} + \overleftarrow{\partial}_{x_2} + \overleftarrow{\partial}_{x_3}) \wedge \overrightarrow{\partial}_x} \dot{\zeta} e^{-\frac{i}{2} (\overleftarrow{\partial}_{x_1} + \overleftarrow{\partial}_{x_2} + \overleftarrow{\partial}_{x_3}) \wedge \overrightarrow{\partial}_x} \dot{\zeta} e^{-\frac{i}{2} (\overleftarrow{\partial}_{x_1} + \overleftarrow{\partial}_{x_2} + \overleftarrow{\partial}_{x_3}) \wedge \overrightarrow{\partial}_x} \partial^{-2} \dot{\zeta} | 0 \rangle \end{aligned} \quad (57)$$

$$(58)$$

The above equation in Fourier space becomes

$$\begin{aligned}
(a) &= -4i\epsilon^2 \int dt' a^5 H \int d^3x \int \prod_{i=1}^6 \frac{d^3k_i}{k_6^2 (2\pi)^{18}} e^{i(\vec{k}_1 \cdot \vec{x}_1 + \vec{k}_2 \cdot \vec{x}_2 + \vec{k}_3 \cdot \vec{x}_3)} \\
&\times \langle 0 | \zeta \left(\vec{k}_1, t_1 + \frac{\vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_3 + \vec{\theta}^0 \cdot \vec{k}_4 + \vec{\theta}^0 \cdot \vec{k}_5 + \vec{\theta}^0 \cdot \vec{k}_6}{2} \right) \\
&\times \langle 0 | \zeta \left(\vec{k}_1, t_1 + \frac{-\vec{\theta}^0 \cdot \vec{k}_1 + \vec{\theta}^0 \cdot \vec{k}_3 + \vec{\theta}^0 \cdot \vec{k}_4 + \vec{\theta}^0 \cdot \vec{k}_5 + \vec{\theta}^0 \cdot \vec{k}_6}{2} \right) \\
&\times \langle 0 | \zeta \left(\vec{k}_1, t_1 + \frac{-\vec{\theta}^0 \cdot \vec{k}_1 - \vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_4 + \vec{\theta}^0 \cdot \vec{k}_5 + \vec{\theta}^0 \cdot \vec{k}_6}{2} \right) \\
&\times \dot{\zeta} \left(\vec{k}_4, t' - \frac{\vec{\theta}^0 \cdot \vec{k}_1 + \vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_3}{2} \right) \\
&\times \dot{\zeta} \left(\vec{k}_5, t' - \frac{\vec{\theta}^0 \cdot \vec{k}_1 + \vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_3}{2} \right) \\
&\times \dot{\zeta} \left(\vec{k}_6, t' - \frac{\vec{\theta}^0 \cdot \vec{k}_1 + \vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_3}{2} \right) | 0 \rangle \\
&\times e^{i(\vec{k}_4 \cdot \vec{x} + \vec{k}_5 \cdot \vec{x} + \vec{k}_6 \cdot \vec{x})} e^{\frac{i}{2}\mathcal{P}}
\end{aligned} \tag{59}$$

Here

$$\mathcal{P} = \left(\vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3 + \left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 \right) \left(\vec{k}_4 + \vec{k}_5 + \vec{k}_6 \right) \right) \tag{60}$$

Since we will express the three-point correlation function in momentum space, we can take the Fourier transform on both side of Eq. (55) and take the limit $t_1 = t_2 = t_3 = t$ to get

$$\begin{aligned}
(a) &= -i \int_{t_0}^t dt' \left(\langle 0 | \zeta_\theta(\vec{k}_1, t) \zeta_\theta(\vec{k}_2, t) \zeta_\theta(\vec{k}_3, t) \mathcal{H}(\sqcup') \right) \\
&= -4i\epsilon^2 \int dt' a^5 H \int d^3x \int \prod_{i=4}^6 \frac{d^3k_i}{k_6^2 (2\pi)^9} \langle 0 | \zeta(\vec{k}_1, t_1) \zeta(\vec{k}_2, t_2) \zeta(\vec{k}_3, t_3) \dot{\zeta}(\vec{k}_4, t_4) \dot{\zeta}(\vec{k}_5, t_5) \dot{\zeta}(\vec{k}_6, t_6) | 0 \rangle \\
&\times e^{i(\vec{k}_4 \cdot \vec{x} + \vec{k}_5 \cdot \vec{x} + \vec{k}_6 \cdot \vec{x})} e^{\frac{i}{2}\mathcal{P}}
\end{aligned} \tag{61}$$

where

$$\begin{aligned}
t_1 &= t + \frac{\vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_3 + \vec{\theta}^0 \cdot \vec{k}_4 + \vec{\theta}^0 \cdot \vec{k}_5 + \vec{\theta}^0 \cdot \vec{k}_6}{2} \\
t_2 &= t + \frac{-\vec{\theta}^0 \cdot \vec{k}_1 + \vec{\theta}^0 \cdot \vec{k}_3 + \vec{\theta}^0 \cdot \vec{k}_4 + \vec{\theta}^0 \cdot \vec{k}_5 + \vec{\theta}^0 \cdot \vec{k}_6}{2} \\
t_3 &= t + \frac{-\vec{\theta}^0 \cdot \vec{k}_1 - \vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_4 + \vec{\theta}^0 \cdot \vec{k}_5 + \vec{\theta}^0 \cdot \vec{k}_6}{2} \\
t_4 &= t' - \frac{\vec{\theta}^0 \cdot \vec{k}_1 + \vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_3}{2} \\
t_5 &= t' - \frac{\vec{\theta}^0 \cdot \vec{k}_1 + \vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_3}{2} \\
t_6 &= t' - \frac{\vec{\theta}^0 \cdot \vec{k}_1 + \vec{\theta}^0 \cdot \vec{k}_2 + \vec{\theta}^0 \cdot \vec{k}_3}{2}
\end{aligned} \tag{62}$$

$$\tag{63}$$

A detailed calculation of this term is presented in Appendix and it is given as (A.11)

$$(a) = \epsilon(2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \frac{e^{\frac{5H\vec{\theta}^0 \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}} e^{\frac{i}{2}(\vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3)}}{K} (k_1^2 k_2^2 + perm.) \tag{64}$$

Here $K = k_1 + k_2 + k_3$. Now similar calculations can be done for the second term in the three-point function (55). Let us represent it as (b),

$$(b) = i \int_{t_0}^t dt' \langle 0 | \mathcal{H}(t') \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) | 0 \rangle \tag{65}$$

The contribution due to this term in momentum space is given by (A.16)

$$(b) = \epsilon(2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \frac{e^{-\frac{5H\vec{\theta}^0 \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}} e^{\frac{i}{2}(\vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3)}}{K} (k_1^2 k_2^2 + perm.)$$

So the contribution to the three-point function of ζ due to the first term of (51) in Fourier space is given as

$$\begin{aligned}
\langle \zeta_\theta(\vec{k}_1, t) \zeta_\theta(\vec{k}_2, t) \zeta_\theta(\vec{k}_3, t) \rangle &= 2\epsilon(2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \\
&\times \frac{\cosh \frac{5H\vec{\theta}^0 \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}}{K} \times e^{\frac{i}{2}(\vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3)} (k_1^2 k_2^2 + perm.)
\end{aligned} \tag{66}$$

This concludes the calculations of the three-point function of the redefined field ζ_n . Now to get the final three-point function of the field ζ we need to consider the second and third

term of equation (51) coming due to field redefinitions. The contribution to the three-point function due to first of these terms can be obtained using Wick's theorem and is given as

$$\langle \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) \rangle = \frac{\eta - \epsilon}{4} (\langle \zeta_\theta(x_1) \zeta_\theta(x_2) \rangle \langle \zeta_\theta(x_1) \zeta_\theta(x_3) \rangle + \text{perm.}) \quad (67)$$

Now

$$\langle \zeta_\theta(x_1) \zeta_\theta(x_2) \rangle = \int \frac{d^3 k_2}{(2\pi)^3} \frac{H^2}{4\epsilon} \frac{1}{k_2^3} e^{-H\vec{\theta}^0 \cdot \vec{k}_2} e^{i\vec{k}_2 \cdot (\vec{x}_1 - \vec{x}_2)} \quad (68)$$

So

$$\begin{aligned} \langle \zeta_\theta(x_1) \zeta_\theta(x_2) \rangle \langle \zeta_\theta(x_1) \zeta_\theta(x_3) \rangle &= \int \frac{d^3 k_2 d^3 k_3}{(2\pi)^9} \frac{H^4}{16\epsilon^2} \frac{1}{k_2^3 k_3^3} e^{-H\vec{\theta}^0 \cdot (\vec{k}_2 + \vec{k}_3)} e^{i(\vec{k}_2 + \vec{k}_3) \cdot \vec{x}_1 - i\vec{k}_2 \cdot \vec{x}_2 - i\vec{k}_3 \cdot \vec{x}_3} \\ &= (2\pi)^3 \int \frac{d^3 k_1 d^3 k_2 d^3 k_3}{(2\pi)^9} \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \frac{1}{k_2^3 k_3^3} \\ &\quad \times e^{H\vec{\theta}^0 \cdot \vec{k}_1} e^{-i\vec{k}_1 \cdot \vec{x}_1 - i\vec{k}_2 \cdot \vec{x}_2 - i\vec{k}_3 \cdot \vec{x}_3} \end{aligned} \quad (69)$$

Here in the second step we have introduced a δ function with integral over k_1 so that it matches with the results of the rest of the terms. So the contribution to the three-point function due to first field redefinition term in momentum space will be

$$\langle \zeta_\theta(\vec{k}_1, t) \zeta_\theta(\vec{k}_2, t) \zeta_\theta(\vec{k}_3, t) \rangle = \frac{\eta - \epsilon}{2} (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \left(\sum_i k_i^3 e^{H\vec{\theta}^0 \cdot \vec{k}_i} \right) \quad (70)$$

Now consider the second field redefinition term in Eq. (51) the contribution due to that is given as

$$\langle \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) \rangle = \frac{\epsilon}{2} (\partial_{x_1}^{-2} (\langle \zeta_\theta(x_1) \zeta_\theta(x_2) \rangle \partial_{x_1}^2 \langle \zeta_\theta(x_1) \zeta_\theta(x_3) \rangle) + \text{perm.}) \quad (71)$$

Now

$$\partial_{x_1}^2 \langle \zeta_\theta(x_1) \zeta_\theta(x_3) \rangle = - \int \frac{d^3 k_3}{(2\pi)^3} \frac{H^2}{4\epsilon} \frac{1}{k_3} e^{-H\vec{\theta}^0 \cdot \vec{k}_3} e^{i\vec{k}_3 \cdot (\vec{x}_1 - \vec{x}_3)} \quad (72)$$

So

$$\begin{aligned} \langle \zeta_\theta(x_1) \zeta_\theta(x_2) \rangle \partial_{x_1}^2 \langle \zeta_\theta(x_1) \zeta_\theta(x_3) \rangle &= - \int \frac{d^3 k_2 d^3 k_3}{(2\pi)^6} \frac{H^4}{16\epsilon^2} \frac{1}{k_2^3 k_3^3} e^{-H\vec{\theta}^0 \cdot (\vec{k}_2 + \vec{k}_3)} e^{i(\vec{k}_2 + \vec{k}_3) \cdot \vec{x}_1 - i\vec{k}_2 \cdot \vec{x}_2 - i\vec{k}_3 \cdot \vec{x}_3} \\ &= -(2\pi)^3 \int \frac{d^3 k_1 d^3 k_2 d^3 k_3}{(2\pi)^9} \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \frac{1}{k_2^3 k_3^3} \\ &\quad \times e^{H\vec{\theta}^0 \cdot \vec{k}_1} e^{-i\vec{k}_1 \cdot \vec{x}_1 - i\vec{k}_2 \cdot \vec{x}_2 - i\vec{k}_3 \cdot \vec{x}_3} \end{aligned} \quad (73)$$

So

$$\begin{aligned} \partial_{x_1}^{-2} (\langle \zeta_\theta(x_1) \zeta_\theta(x_2) \rangle \partial_{x_1}^2 \langle \zeta_\theta(x_1) \zeta_\theta(x_3) \rangle) &= (2\pi)^3 \int \frac{d^3 k_1 d^3 k_2 d^3 k_3}{(2\pi)^9} \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \frac{1}{k_1^3 k_2^3 k_3^3} \\ &\quad \times e^{H\vec{\theta}^0 \cdot \vec{k}_1} e^{-i\vec{k}_1 \cdot \vec{x}_1 - i\vec{k}_2 \cdot \vec{x}_2 - i\vec{k}_3 \cdot \vec{x}_3} \end{aligned} \quad (74)$$

So the contribution due to this term in Fourier space will be

$$\langle \zeta_\theta(\vec{k}_1, t) \zeta_\theta(\vec{k}_2, t) \zeta_\theta(\vec{k}_3, t) \rangle = \frac{\epsilon}{2} (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \left(\sum_{i \neq j} k_i k_j^2 e^{H\vec{\theta}^\partial \cdot \vec{k}_i} \right) \quad (75)$$

Now combining all the results from (66), (70), (75) for the various contributions to the three-point function of ζ , we get the final three-point function using Eq. (51) in momentum space as

$$\langle \zeta_\theta(\vec{k}_1, t) \zeta_\theta(\vec{k}_2, t) \zeta_\theta(\vec{k}_3, t) \rangle = (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \mathcal{A} \quad (76)$$

Where

$$\begin{aligned} \mathcal{A} = & 4\epsilon \frac{\cosh \frac{5H\vec{\theta}^\partial \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2} \times e^{\frac{i}{2}(\vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3)}}{K} \left(\sum_{i < j} k_i^2 k_j^2 \right) + \frac{\eta - \epsilon}{2} \left(\sum_i k_i^3 e^{H\vec{\theta}^\partial \cdot \vec{k}_i} \right) \\ & + \frac{\epsilon}{2} \left(\sum_{i \neq j} k_i k_j^2 e^{H\vec{\theta}^\partial \cdot \vec{k}_i} \right) \end{aligned} \quad (77)$$

This is the main result of this paper. In the limit $\theta^{\mu\nu} \rightarrow 0$ the above expression becomes similar to expression for the three-point function in commutative spacetime (Eq. (4.5) and (4.6) of Maldacena [10]). Now due to translational invariance $\vec{k}_1 + \vec{k}_2 + \vec{k}_3 = 0$ and on comparing our results with the commutative case [10, 18] we see that the first term in (77) is modified due to a phase factor that depends on θ_{ij} , while the second and the last terms are modified by exponential factors. The three-point correlation function here is complex so to see its observational effects we again take its self adjoint given as [16]

$$\begin{aligned} \langle \zeta_\theta(\vec{k}_1, t) \zeta_\theta(\vec{k}_2, t) \zeta_\theta(\vec{k}_3, t) \rangle_M &= \frac{1}{2} \left(\langle \zeta_\theta(\vec{k}_1, t) \zeta_\theta(\vec{k}_2, t) \zeta_\theta(\vec{k}_3, t) \rangle + \langle \zeta_\theta(-\vec{k}_1, t) \zeta_\theta(-\vec{k}_2, t) \zeta_\theta(-\vec{k}_3, t) \rangle \right) \\ &= (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \left[\frac{4\epsilon \cos\left(\frac{\vec{k}_1 \wedge \vec{k}_2}{2}\right)}{K} \left(\sum_{i < j} k_i^2 k_j^2 \right) \right. \\ &\quad \left. + \frac{\eta - \epsilon}{2} \left(\sum_i k_i^3 \cosh(H\vec{\theta}^\partial \cdot \vec{k}_i) \right) + \frac{\epsilon}{2} \left(\sum_{i \neq j} k_i k_j^2 \cosh(H\vec{\theta}^\partial \cdot \vec{k}_i) \right) \right] \end{aligned} \quad (78)$$

4. IMPLICATIONS FOR OBSERVATIONS

The non-gaussianity in CMB is described in terms of the angular three-point correlation functions in harmonic space called as "angular bispectrum", which is related to the three-dimensional bispectrum of the primordial curvature perturbations defined as [24, 25]

$$\langle \zeta(\vec{k}_1, t) \zeta(\vec{k}_2, t) \zeta(\vec{k}_3, t) \rangle = (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) B_\zeta(k_1, k_2, k_3) \quad (79)$$

We can generalize the above definition of bispectrum for the twisted quantum fields in noncommutative space time and it can be expressed using (78) as

$$B_{\zeta_\theta}(\vec{k}_1, \vec{k}_2, \vec{k}_3) = \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \left[\frac{4\epsilon \cos\left(\frac{(\vec{k}_1 \wedge \vec{k}_2)}{2}\right)}{K} \left(\sum_{i<j} k_i^2 k_j^2 \right) + \frac{\eta - \epsilon}{2} \left(\sum_i k_i^3 \cosh\left(H\vec{\theta}^0 \cdot \vec{k}_i\right) \right) \right. \\ \left. + \frac{\epsilon}{2} \left(\sum_{i \neq j} k_i k_j^2 \cosh\left(H\vec{\theta}^0 \cdot \vec{k}_i\right) \right) \right] \quad (80)$$

Here the bispectrum also breaks the statistical isotropy. The anisotropic bispectrum also arises in the cases where the vector fields are also present during inflation [25]. In [26] the method to analyze these models in the light of new CMB data is derived. Current observational limits on non-gaussianity are given in terms of a non-linearity parameter f_{NL} that determines the amplitude and scale dependence of non-gaussianity. We define f_{NL} in a similar way as [25] where it is assumed that the corrections to the standard power spectrum due to statistical anisotropy are very small. So

$$f_{NL} = \frac{5}{6} \frac{B_{\zeta_\theta}(\vec{k}_1, \vec{k}_2, \vec{k}_3)}{P_\zeta(k_1)P_{\zeta_\theta}(k_2) + P_\zeta(k_2)P_\zeta(k_3) + P_\zeta(k_1)P_\zeta(k_3)} \quad (81)$$

Using the power spectrum (35) it becomes

$$f_{NL} = \frac{5}{6} \frac{1}{\sum_i k_i^3} \left[4\epsilon \frac{\cos\left(\frac{\vec{k}_1 \wedge \vec{k}_2}{2}\right)}{K} \sum_{i<j} (k_i^2 k_j^2) + \frac{\eta - \epsilon}{2} \left(\sum_i k_i^3 \cosh\left(H\vec{\theta}^0 \cdot \vec{k}_i\right) \right) \right. \\ \left. + \frac{\epsilon}{2} \left(\sum_{i \neq j} k_i k_j^2 \cosh\left(H\vec{\theta}^0 \cdot \vec{k}_i\right) \right) \right] \quad (82)$$

This kind of f_{NL} generally arises where the curvature perturbation is expressed as $\zeta_g = \zeta_g + \frac{3}{5}\zeta_g^2$ and f_{NL} peaks at the so called squeezed triangle limit defined as $|\vec{k}_1| = |\vec{k}_2| = k$ and $|\vec{k}_3| \ll k$. So in a similar fashion the f_{NL} for noncommutative case in the above limit is given as

$$f_{NL} = \frac{5}{12} \left[2\epsilon \cos\left(\frac{\vec{k}_1 \wedge \vec{k}_2}{2}\right) + \frac{\eta}{2} \left(\cosh\left(H\vec{\theta}^0 \cdot \vec{k}_1\right) + \cosh\left(H\vec{\theta}^0 \cdot \vec{k}_2\right) \right) \right] \quad (83)$$

It is clear from the above expression that the amplitude of f_{NL} is very small and of the order of slow-roll parameters for the case of small statistical anisotropy. But it has scale dependence and direction dependence that can help us to distinguish it from the commutative case. The current limits on the amplitude of f_{NL} are [27] $f_{NL} = 48 \pm 20$ for squeezed triangle limit at 68% confidence level and the Planck is supposed to see the $f_{NL} \sim 5$. The running of the non-gaussianity n_G for $\vec{\theta}^0 = 0$ in our case is similar to [16], and it is shown in [28, 29] that we will be able to constraint n_{NG} with a $1 - \sigma$ uncertainty of $\Delta n_{NG} \sim 0.1$ with ongoing Planck and future CMB and large scale structure observations. Hence even though the amplitude of f_{NL} with the noncommutative geometry is of the order of slow-roll parameters, there is a possibility to see the effects of scale dependence of f_{NL} due to noncommutativity in future observations.

5. CONCLUSIONS

Detection of primordial non-gaussianity in the CMB anisotropy and large scale structure is the main challenge of current and future observations and it can play an important role in discriminating various models of inflation. In this paper we have calculated the primordial non-gaussianity in single field inflation with spacetime noncommutativity. We have used Maldacena's approach [10] to compute the two-point and three-point correlation functions for the comoving curvature perturbation ζ for the noncommutative case described by [12]. Both the power spectrum and the bispectrum for this model are direction dependent and breaks the statistical isotropy due to the preferred direction of $\hat{\theta}$. This direction dependent power spectrum was analyzed by [13] to put constraints on the scale of noncommutativity in the light of WMAP5 data and it was concluded that the WMAP5 data at high l is not sufficient to constraint the noncommutative scale θ . Since Planck will measure the CMB anisotropy at high-precession up to $l \geq 2500$, the author and collaborators plan to test the power spectrum (31) after the release of Planck data. The statistical anisotropic bispectrum can be extracted from the three-point correlation function of CMB [26] and for $f_{NL} \approx 30$, future experiments could be sensitive to a ratio of the anisotropic to the isotropic amplitudes of the bispectrum up to 10%. The amplitude of the non-linearity parameter f_{NL} for our case is very small for small statistical anisotropy but it has a high scale dependence. Ongoing Planck and future CMB and large scale structure observations would be able to measure the running of non-gaussianity up-to $1 - \sigma$ uncertainty of $\Delta n_{NG} \sim 0.1$ [29], that can help us in distinguishing the effects due to noncommutativity. With the help of these observational signatures of noncommutative inflation we will, in future, be able to determine the scale of noncommutativity.

6. ACKNOWLEDGEMENT

We thank A. P. Balachandran for very helpful discussions and valuable suggestions.

Appendix

The integral appearing in the calculation of the first term in the three-point function (55) in Fourier space can be read from Eq. (61) as

$$\begin{aligned}
 (a) = & -4i\epsilon^2 \int dt' a^5 H \int d^3x \int \prod_{i=4}^6 \frac{d^3k_i}{k_6^2 (2\pi)^9} \langle 0 | \zeta(\vec{k}_1, t_1) \zeta(\vec{k}_2, t_2) \zeta(\vec{k}_3, t_3) \dot{\zeta}(\vec{k}_4, t_4) \dot{\zeta}(\vec{k}_5, t_5) \dot{\zeta}(\vec{k}_6, t_6) | 0 \rangle \\
 & \times e^{i(\vec{k}_4 \cdot \vec{x} + \vec{k}_5 \cdot \vec{x} + \vec{k}_6 \cdot \vec{x})} e^{\frac{i}{2}\mathcal{P}},
 \end{aligned} \tag{A.1}$$

where t_i s are given by Eq. (62). Now we will calculate the six-point function entering in the above integrand separately and denote it as A . So

$$A = \langle 0 | \zeta(\vec{k}_1, t_1) \zeta(\vec{k}_2, t_2) \zeta(\vec{k}_3, t_3) \dot{\zeta}(\vec{k}_4, t_4) \dot{\zeta}(\vec{k}_5, t_5) \dot{\zeta}(\vec{k}_6, t_6) | 0 \rangle \tag{A.2}$$

Now using Wick's theorem and leaving the disconnected diagrams we will get 6 terms in above expression. Let us consider one of them and denote it by A_1 so

$$A_1 = \langle 0 | \left[\zeta^+(\vec{k}_1, t_1), \dot{\zeta}^-(\vec{k}_4, t_4) \right] \left[\zeta^+(\vec{k}_2, t_2), \dot{\zeta}^-(\vec{k}_5, t_5) \right] \left[\zeta^+(\vec{k}_3, t_3), \dot{\zeta}^-(\vec{k}_6, t_6) \right] | 0 \rangle \quad (\text{A.3})$$

where the ζ^+ and ζ^- denote the positive and negative frequency part of the quantum field ζ (see (24)). Now since

$$\left[\zeta^+(\vec{k}_1, t_1), \dot{\zeta}^-(\vec{k}_4, t_4) \right] = (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_4) u(\vec{k}_1, t_1) \dot{u}^*(-\vec{k}_4, t_4) \quad (\text{A.4})$$

we have

$$\begin{aligned} A_1 &= (2\pi)^9 \delta^3(\vec{k}_1 + \vec{k}_4) \delta^3(\vec{k}_2 + \vec{k}_5) \delta^3(\vec{k}_3 + \vec{k}_6) u(\vec{k}_1, t_1) u(\vec{k}_2, t_2) u(\vec{k}_3, t_3) \\ &\quad \times \dot{u}^*(-\vec{k}_4, t_4) \dot{u}^*(-\vec{k}_5, t_5) \dot{u}^*(-\vec{k}_6, t_6) \end{aligned} \quad (\text{A.5})$$

Putting this back to the integral (59) and denoting the contribution due to this term as $(a)_1$ and doing the delta integrals we get

$$\begin{aligned} (a)_1 &= -4i\epsilon^2 \int_{t_0}^t dt' a^5 H \int d^3x \frac{1}{k_3^2} u(\vec{k}_1, t_1) u(\vec{k}_2, t_2) u(\vec{k}_3, t_3) \dot{u}^*(\vec{k}_1, t_4) \dot{u}^*(\vec{k}_2, t_5) \dot{u}^*(\vec{k}_3, t_6) \\ &\quad \times e^{i(\vec{k}_4 \cdot \vec{x} + \vec{k}_5 \cdot \vec{x} + \vec{k}_6 \cdot \vec{x})} e^{\frac{i}{2}\mathcal{P}_1} \\ &= -4i\epsilon^2 (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) \int_{t_0}^t dt' a^5 H \frac{1}{k_3^2} u(\vec{k}_1, t_1) u(\vec{k}_2, t_2) u(\vec{k}_3, t_3) \\ &\quad \times \dot{u}^*(\vec{k}_1, t_4) \dot{u}^*(\vec{k}_2, t_5) \dot{u}^*(\vec{k}_3, t_6) e^{\frac{i}{2}\mathcal{P}_1} \end{aligned} \quad (\text{A.6})$$

where $\mathcal{P}_1 = \mathcal{P}|_{\vec{k}_4=-\vec{k}_1, \vec{k}_5=-\vec{k}_2, \vec{k}_6=-\vec{k}_3}$ and t_i s are also calculated using these values of momenta. Here the limit of integration goes from $t_0 = -\infty$ to $t = \infty$. To solve the integral we will go to conformal time where the above limits correspond to $\tau \rightarrow (-\infty, 0)$. Now from Eq. (28) we have

$$u(\vec{k}, \tau) = \frac{v_{\vec{k}}}{z} = \frac{iH}{\sqrt{4\epsilon k^3}} (1 + ik\tau) e^{-ik\tau} \quad (\text{A.7})$$

Since from Eq. (40) we know that the conformal time corresponding to t_i s will be $\tau \times e^{\theta \text{ dependent term}}$ and from Eq. (62) we have terms like $(t + \theta \text{ dependent term})$ for t_1, t_2, t_3 so for $t \rightarrow \infty$ or $\tau \rightarrow 0$ conformal time corresponding to t_1, t_2, t_3 will be zero. So in conformal time $u(\vec{k}_1, t_1) u(\vec{k}_2, t_2) u(\vec{k}_3, t_3) \rightarrow u(\vec{k}_1, 0) u(\vec{k}_2, 0) u(\vec{k}_3, 0)$. Now we denote the conformal time corresponding to t' by τ so $\dot{u}^*(\vec{k}_1, t_4) \rightarrow \frac{1}{a} \frac{du^*(\vec{k}_1, \tau_4)}{d\tau}$ and from (40) and (62) we get

$\tau_4 = \tau_5 = \tau_6 = \tau e^{\frac{H\vec{\theta} \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}}$. Now

$$\frac{du^*(\vec{k}_1, \tau_4)}{d\tau} = \frac{-iH}{\sqrt{4\epsilon k^3}} k_1^2 \tau e^{H\vec{\theta} \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)} e^{ik_1 \tau e^{\frac{H\vec{\theta} \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}}} \quad (\text{A.8})$$

So the integral (A.6) becomes

$$\begin{aligned}
(a)_1 &= -i\epsilon(2\pi)^3\delta^3\left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3\right) \frac{H^7}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \\
&\times \int_{-\infty}^0 a^3 \tau^3 k_1^2 k_2^2 e^{\frac{i}{2}\mathcal{P}_1} e^{3H\vec{\theta}^0 \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)} e^{iK\tau e^{\frac{H\vec{\theta}^0 \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}}} \\
&= \epsilon(2\pi)^3\delta^3\left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3\right) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \frac{k_1^2 k_2^2}{K} e^{\frac{i}{2}\mathcal{P}_1} e^{\frac{5H\vec{\theta}^0 \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}}
\end{aligned} \tag{A.9}$$

Here we have rotated the contour from $(-\infty, 0)$ to $i(\infty, 0)$ and $K = k_1 + k_2 + k_3$. Now to calculate $\mathcal{P}_1$ let us recall (60)

$$\mathcal{P} = \vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3 + \vec{k}_1 \wedge \left(\vec{k}_4 + \vec{k}_5 + \vec{k}_6\right) + \vec{k}_2 \wedge \left(\vec{k}_4 + \vec{k}_5 + \vec{k}_6\right) + \vec{k}_3 \wedge \left(\vec{k}_4 + \vec{k}_5 + \vec{k}_6\right) \tag{A.10}$$

Hence $\mathcal{P}_1 = \vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3$. Now rest of the terms in (A.2) can be found by different permutations of k_4, k_5, k_6 and the phase factors will be same as $\mathcal{P}_1$ after imposing the different conditions due to delta function integrals. So from equation (61) we get the first term of the right hand side of equation (55) in Fourier space as

$$(a) = \epsilon(2\pi)^3\delta^3\left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3\right) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \frac{e^{\frac{5H\vec{\theta}^0 \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}} e^{\frac{i}{2}(\vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3)}}{K} (k_1^2 k_2^2 + perm.) \tag{A.11}$$

Now the second term in the three-point function (55) is denoted as (b) and can be read from Eq. (A.13) as

$$\begin{aligned}
(b) &= i \int_{t_0}^t dt' \langle 0 | \mathcal{H}(t') \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) | 0 \rangle \\
&= -4i\epsilon^2 \int dt' a^5 H \int d^3x \langle 0 | \dot{\zeta}^2 \partial^{-2} \dot{\zeta} \Big|_{t', \vec{x}} e^{\frac{i}{2} \overleftarrow{\partial}_{x\mu} \wedge P_\nu} \zeta_\theta(x_1) \zeta_\theta(x_2) \zeta_\theta(x_3) | 0 \rangle \\
&= -4i\epsilon^2 \int dt' a^5 H \int d^3x \langle 0 | \dot{\zeta}^2 \partial^{-2} \dot{\zeta} \Big|_{t', \vec{x}} e^{-\frac{i}{2} \overleftarrow{\partial}_x \wedge (\overrightarrow{\partial}_{x_1} + \overrightarrow{\partial}_{x_2} + \overrightarrow{\partial}_{x_3})} \\
&\times \zeta(x_1) \zeta(x_2) \zeta(x_3) | 0 \rangle e^{-\frac{i}{2} (\overleftarrow{\partial}_{x_1} \wedge \overleftarrow{\partial}_{x_2} + \overleftarrow{\partial}_{x_2} \wedge \overleftarrow{\partial}_{x_3} + \overleftarrow{\partial}_{x_1} \wedge \overleftarrow{\partial}_{x_3})}
\end{aligned} \tag{A.12}$$

Here we have used (36). Now in the Fourier space we get

$$\begin{aligned}
(b) &= -4i\epsilon^2 \int dt' a^5 H \int d^3x \int \prod_{i=1}^6 \frac{d^3k}{-k_6^2 (2\pi)^{18}} e^{i(\vec{k}_4 + \vec{k}_5 + \vec{k}_6) \cdot \vec{x}} e^{-\frac{i}{2} \overleftarrow{\partial}_x \wedge (\overrightarrow{\partial}_{x_1} + \overrightarrow{\partial}_{x_2} + \overrightarrow{\partial}_{x_3})} \\
&\times \langle 0 | \dot{\zeta}(\vec{k}_4, t') \dot{\zeta}(\vec{k}_5, t') \dot{\zeta}(\vec{k}_6, t') \zeta(\vec{k}_1, t_1) \zeta(\vec{k}_2, t_2) \zeta(\vec{k}_3, t_3) | 0 \rangle \\
&\times e^{i(\vec{k}_1 \cdot \vec{x}_1 + \vec{k}_2 \cdot \vec{x}_2 + \vec{k}_3 \cdot \vec{x}_3)} e^{-\frac{i}{2} (\overleftarrow{\partial}_{x_1} \wedge \overleftarrow{\partial}_{x_2} + \overleftarrow{\partial}_{x_2} \wedge \overleftarrow{\partial}_{x_3} + \overleftarrow{\partial}_{x_1} \wedge \overleftarrow{\partial}_{x_3})} \\
&= 4i\epsilon^2 \int dt' a^5 H \int d^3x \int \prod_{i=1}^6 \frac{d^3k}{k_6^2 (2\pi)^{18}} e^{i(\vec{k}_4 + \vec{k}_5 + \vec{k}_6) \cdot \vec{x}} e^{i(\vec{k}_1 \cdot \vec{x}_1 + \vec{k}_2 \cdot \vec{x}_2 + \vec{k}_3 \cdot \vec{x}_3)} \\
&\times \langle 0 | \dot{\zeta}(\vec{k}_4, t_4) \dot{\zeta}(\vec{k}_5, t_5) \dot{\zeta}(\vec{k}_6, t_6) \zeta(\vec{k}_1, t_1) \zeta(\vec{k}_2, t_2) \zeta(\vec{k}_3, t_3) | 0 \rangle e^{\frac{i}{2} \vec{P}}
\end{aligned} \tag{A.13}$$

Here

$$\begin{aligned}
\tilde{\mathcal{P}} &= \vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3 - \left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 \right) \wedge \left(\vec{k}_4 + \vec{k}_5 + \vec{k}_6 \right) \\
t_1 &= t + \frac{\vec{\theta}^0 \cdot \left(\vec{k}_2 + \vec{k}_3 - \vec{k}_4 - \vec{k}_5 - \vec{k}_6 \right)}{2} \\
t_2 &= t + \frac{\vec{\theta}^0 \cdot \left(-\vec{k}_1 + \vec{k}_3 - \vec{k}_4 - \vec{k}_5 - \vec{k}_6 \right)}{2} \\
t_3 &= t + \frac{\vec{\theta}^0 \cdot \left(\vec{k}_1 - \vec{k}_2 - \vec{k}_4 - \vec{k}_5 - \vec{k}_6 \right)}{2} \\
t_4 &= t' + \frac{\vec{\theta}^0 \cdot \left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 \right)}{2} \\
t_5 &= t' + \frac{\vec{\theta}^0 \cdot \left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 \right)}{2} \\
t_6 &= t' + \frac{\vec{\theta}^0 \cdot \left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 \right)}{2}
\end{aligned} \tag{A.14}$$

$$\tag{A.15}$$

Now all the calculations can be done for (b) as earlier and the final answer is

$$(b) = \epsilon(2\pi)^3 \delta^3 \left(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 \right) \frac{H^4}{16\epsilon^2} \prod_{i=1}^3 \frac{1}{k_i^3} \frac{e^{-\frac{5H\vec{\theta}^0 \cdot (\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{2}} e^{\frac{i}{2}(\vec{k}_1 \wedge \vec{k}_2 + \vec{k}_2 \wedge \vec{k}_3 + \vec{k}_1 \wedge \vec{k}_3)}}{K} (k_1^2 k_2^2 + perm.) \tag{A.16}$$

-
- [1] A. H. Guth, Phys. Rev. **D23**, 347 (1981).
 - [2] G. F. Smoot, C. L. Bennett, A. Kogut, E. L. Wright, J. Aymon, N. W. Boggess, E. S. Cheng, G. De Amici *et al.*, Astrophys. J. **396**, L1-L5 (1992).
 - [3] D. N. Spergel *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **148**, 175-194 (2003). [astro-ph/0302209]
 - [4] E. Komatsu *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **192**, 18 (2011). [arXiv:1001.4538 [astro-ph.CO]].
 - [5] G. Hinshaw, D. Larson, E. Komatsu, D. N. Spergel, C. L. Bennett, J. Dunkley, M. R.olta and M. Halpern *et al.*, arXiv:1212.5226 [astro-ph.CO].
 - [6] [Planck Collaboration], [astro-ph/0604069] <http://www.rssd.esa.int/Planck>.
 - [7] D. Baumann *et al.* [CMBPol Study Team Collaboration], AIP Conf. Proc. **1141**, 10 (2009) [arXiv:0811.3919 [astro-ph]].
 - [8] L. Amendola *et al.* [Euclid Theory Working Group Collaboration], arXiv:1206.1225 [astro-ph.CO].
 - [9] E. Komatsu, astro-ph/0206039.
 - [10] J. M. Maldacena, JHEP **0305**, 013 (2003) [astro-ph/0210603].

- [11] E. Akofor, A. P. Balachandran and A. Joseph, Int. J. Mod. Phys. A **23**, 1637 (2008) [arXiv:0803.4351 [hep-th]].
- [12] E. Akofor, A. P. Balachandran, S. G. Jo, A. Joseph and B. A. Qureshi, JHEP **0805**, 092 (2008) [arXiv:0710.5897 [astro-ph]].
- [13] E. Akofor, A. P. Balachandran, A. Joseph, L. Pekowsky and B. A. Qureshi, Phys. Rev. D **79**, 063004 (2009) [arXiv:0806.2458 [astro-ph]].
- [14] C. -S. Chu, B. R. Greene and G. Shiu, Mod. Phys. Lett. A **16**, 2231 (2001) [hep-th/0011241].
- [15] R. Brandenberger and P. -M. Ho, Phys. Rev. D **66**, 023517 (2002) [AAPPS Bull. **12N1**, 10 (2002)] [hep-th/0203119].
- [16] T. S. Koivisto and D. F. Mota, JHEP **1102**, 061 (2011) [arXiv:1011.2126 [astro-ph.CO]].
- [17] D. H. Lyth and Y. Rodriguez, Phys. Rev. Lett. **95**, 121302 (2005) [astro-ph/0504045].
- [18] D. Seery and J. E. Lidsey, JCAP **0506**, 003 (2005) [astro-ph/0503692].
- [19] V. F. Mukhanov, H. A. Feldman and R. H. Brandenberger, Phys. Rept. **215**, 203 (1992).
- [20] R. L. Arnowitt, S. Deser and C. W. Misner, gr-qc/0405109.
- [21] D. H. Lyth and Y. Rodriguez, Phys. Rev. D **71**, 123508 (2005) [astro-ph/0502578].
- [22] X. Chen, M. -x. Huang, S. Kachru and G. Shiu, JCAP **0701**, 002 (2007) [hep-th/0605045].
- [23] S. Weinberg, Phys. Rev. D **72**, 043514 (2005) [hep-th/0506236].
- [24] E. Komatsu, Class. Quant. Grav. **27**, 124010 (2010) [arXiv:1003.6097 [astro-ph.CO]].
- [25] E. Dimastrogiovanni, N. Bartolo, S. Matarrese and A. Riotto, Adv. Astron. **2010**, 752670 (2010) [arXiv:1001.4049 [astro-ph.CO]].
- [26] N. Bartolo, E. Dimastrogiovanni, M. Liguori, S. Matarrese and A. Riotto, JCAP **1201**, 029 (2012) [arXiv:1107.4304 [astro-ph.CO]].
- [27] J. -Q. Xia, C. Baccigalupi, S. Matarrese, L. Verde and M. Viel, JCAP **1108**, 033 (2011) [arXiv:1104.5015 [astro-ph.CO]].
- [28] E. Sefusatti, M. Liguori, A. P. S. Yadav, M. G. Jackson and E. Pajer, JCAP **0912**, 022 (2009) [arXiv:0906.0232 [astro-ph.CO]].
- [29] A. Becker, D. Huterer and K. Kadota, JCAP **1212**, 034 (2012) [arXiv:1206.6165 [astro-ph.CO]].